

Pedersen M (2005). A review of evidence based research on laryngeal remobilisation for bilateral paralysis: Round table in Rome IFOS

Round table in Rome

A Review of evidence based research on laryngeal remobilisation for bilateral paralysis

Lx 330Deg 262.5Hz 68.3dB



M. Pedersen MD Dr. Med. Sci
ENT surgeon, FRSM
The Medical Centre, Voice Unit
Østergade 18
DK 1100 Copenhagen
Denmark,
www.mpedersen.org

Lx 330 Deg 262.5Hz 68.3dB



Abstract

A scientific search of papers was made for the round table that included the topic of laryngeal remobilisation for bilateral paralysis.

One protocol was found in the Cochrane evidence based library for unilateral recurrent nerve palsy (Lowe DA and Hoare TJ, 2003). But bilateral immobilisation was excluded due to the complex ethiology of the disorder.

This argument is not medically, ethically relevant in relation to treatment.

On that basis a suggestion for a structure for future evidence and an eventual protocol for an evidence based study has been made, the adequate statistical number of patients in the different groups for a reasonable power of the study is not the problem, because evidence based pilot studies also have to be made, the ethical aspects are that without controls the treatments will always be doubted.

Key words: paralysis- bilateral- vocal cords- remobilisation- voice- respiration- communication- quality of life

Introduction

A search was made in the Cochrane Database for evidence based voice studies of laryngeal remobilisation for bilateral paralysis, as it earlier was done in other laryngeal connections (Pedersen M and McGlashan J, 2000, Hopkins C, Yoysaf U, Pedersen M 2004).


None were found using the traditional Cochrane search strategies, one Cochrane protocol was found related to unilateral recurrent nerve palsy, bilateral palsy was not taken into account because of the many sided aspects of ethiology.

The studies found in other connections were mostly related to SURGERY, which was found to give better respiration possibilities, and this surgical approach has been refined over the later years e.g. in the United States at the Mayo Clinic, a certain aggravated voice situation was accepted by patients and surgeons but not documented.

Respiration related quality of life measurements were not found in the literature

Life quality scores were seldom found and without evidence.

Voice related quality of life and objective measure of the voice were seldom given



Material and Method

We have suggested that clinical voice laboratories in the European Union should include a **standard clinical equipment**, an **optimal clinical equipment**, and for **research clinics** also further equipments, the following settings should be required including medical doctors, physicists, statisticians, psychologists and eventually others first introduced by Pedersen M and Benini A (2001):

Lx 330Deg 262.5Hz 68.3dB



Physiological area

Standard equipment:

Stroboscopy, Airflow measurement, Mean Flow Rate, Phonation time, Videokymography

Optimal equipment also:

Electroglottography, Respiratory measurements (long and short time) Air pressure, Electromyography, Articulatory measurements (three-dimensional magnetic sensor)

Lx 330Deg 262.5Hz 68.3dB



Physiological area

Further research

Equipment:

Videostroboscopy with quantitative computing, Instrumentation for brain stem, brain flow and other brain activity measurements, ultra sound, stem cells and genetics. Evidence based approaches

Lx 330Deg 262.5Hz 68.3dB



Acoustical area

Standard equipment:

Recording procedure: Audio tape (analogue recording),
Fundamental Frequency with jitter%, Intensity with shimmer
%, Signal to noise ratio, signal to harmonics ratio, nasality,
spectral analysis (FFT,LTAS power spectrum)

Optimal equipment also:

Recording procedure (digital recording)
Phonetograms (2- or 3- dimensional) for speech, singing and
shouting, Phonation index, Diplophonia, Voice breaks,
Simultaneous video- and Sound recording (for analysis)

Lx 330Deg 262.5Hz 68.3dB



Acoustical area

Further research

Equipment:

Stroboscopy combined with glottography and phonetography. Motor speech profile measurements, evidence based efficiency measurements based upon air pressure, air flow and acoustic signal

Lx 330Deg 262.5Hz 68.3dB



Perceptual and Psychological area:

Standard methods:

Listening tests including the GRBAS test
Other voice quality tests including for nasality
Standard methods for registration of the patients
subjective statements of illness, the Voice Handicap Index
Voice Related Quality of Life (VRQOL).

Optimal methods also:

Objective evidence based speech acoustics related
perception and music acoustics related perception
(Wøldike test of musicality, (Pedersen M)), video-
and acoustic recording (e.g. by stuttering and
spasmodic dysphonia)

Lx:330Deg 262.5Hz 68.3dB



Perceptual and Psychological area:

Further research

Equipment:

Objective registration of moods at the physiological and acoustical Level. Evidence based approaches.

Lx 330Deg 262.5Hz 68.3dB



Results

In the Cochrane library by double search only **one protocol** for surgery for vocal cord paralysis and paresis was found (Lowe DA and Hoare TJ 2003). **Surgery for bilateral vocal cord paralysis was excluded** in the protocol due to considerable differences in the 1. management, 2. surgical techniques and 3. aim of the treatment. **In addition patients with vagal rather than recurrent nerve palsy were excluded** because vagal palsy often is more difficult to treat.

No reviews were found.

In the Cochrane library 4 controlled clinical trials on laryngeal immobilisation of one vocal cord were found, all focusing on the use of EMG in unilateral paralysis. None were found for aspects of bilateral immobilisation.

Due to the *evidence related results of EMG on vocal cords in the Cochrane Library*, the EMG studies are presented, the use of EMG is stressed:


Lx.330Deg 202.5Hz 68.3dB



Mostafa, (2004)

Mostafa BE, Gadallah NA, Nassar NM, AL Ibiary HM, Fahmy HA, Fouada NM. (2004) from Ain-Shams University in Cairo examined 35 patients prospectively with a control group of 10 normals.

This means that a case-control study was made: The specificity of EMG was 100% detecting vocal fold immobility. The sensitivity was 65.7%. 26% had isolated recurrent nerve lesion, 74% combined superior laryngeal nerve and recurrent nerve lesions. After 6 months 40% were recovered, the specificity of EMG was 100% in predicting recovery.



Wahl (1998)

Wahl RA and Rimpl I from Germany in a consecutive series of 1143 first operations for benign nodular goitre with 1928 “nerves at risk” found that the rate of permanent recurrent laryngeal nerve palsy was higher after non-identification of the nerve (0.6% vs 0%).

As a rule the nerve should be identified, especially in conventional subtotal resection, when possible without further mobilizing manipulations.

Lx 330Deg 262.5Hz 68.3dB



Hayashi (1997)

Hayashi M, Isozaki I, Oda M, Tanabe H and Kimura J from Tokio Metropolitan Neurological Hospital examined 6 patients with multiple system atrophy compared with 6 normals.

In multiple system atrophy the small myelinated fibres innervating the vocal cord are affected first without obvious clinical signs.

The patient develops vocal cord palsy **only after loss of the large myelinated fibres** which mostly comprise the alpha motor axons that innervate the intrinsic laryngeal muscles.

Lx 330Deg 262.5Hz 08_3dB



Koch (1996)

Koch B, Boettcher M, Huschitt N and Hülsewede R from Germany prospectively compared two groups of subtotal resections of the thyroid (382/413) with “nerves at risk”,

in each group two patients had transient vocal cord paralysis that recovered within 4 months.

The demand for obligatory intraoperative identification of the recurrent nerve seemed not to be tenable

Lx 330Deg 262.5Hz 68.3dB

A total search of the literature included 194 papers focusing on aspects *for subgroups, non-evidence based of etiology, diagnosis, treatment and swallowing.*

ETHIOLOGY

The papers focused on post goitre operation, trauma, malignancy, virus, neurological, metabolic and toxic disorders.

It can be commented that a very good **GENERAL medical knowledge** of aspects of:

Endocrinology

Malignancy

Trauma

Virus

Neurology

Metabolic disorders

Toxic disorders

is necessary **prophylactically** also in the area of remobilisation of the vocal cords, because with such knowledge many cases can be treated without surgery and eventual remobilisation. The ethical aspects in this area must be taken into account.

A survey of tests to be made is given e.g. by Ernster JA et al.(2002). A differentiation to pre cancer situations is suggested by Arens C, Schöberlein S and Glanz HK (2005)

DIAGNOSIS

included EMG, evaluation of mobilisation of the arytenoids, scars especially in the interarytenoid region, evaluation of decline of respiratory function including spirometry, peak flow, stroboscopy and voice analysis.

(e.g. peak flow, phonation time, mean flow rate and voice intensity were analysed after the operation (Remacle M et al. 1996))

Lx 330Deg 262.5Hz 68.3dB



It can be commented that a good **GENERAL knowledge** with the aspects of:

EMG

Mobilisation of the arytenoids

Scars especially between the arytenoids

Respiration function

Spirometry

Peakflow

Videostroboscopy

Voice analysis

Voice treatment

General health (SF 36)

Subjective voice related complaints

is necessary **prophylactically**, again because with such knowledge many cases can be treated without surgery and eventual remobilisation, taking care of intermediate exacerbations that are not immobilisation related, and which could also happen after remobilisation.

SURGICAL / NONSURGICAL TREATMENT

Focused on indications for tracheostomy, arytenoidectomy, cordectomy, and cordopexy.

The reinnervation techniques and electric pacing are considered to be experimental.

The works by Tucker HM and Ogura JH in animals, and Fex S (1970) conclude that the phrenic nerve implantation in the posticus muscle, for reinnervation of abduction at inspiration is the better choice.

Zelear DL et al. (2003), and others have tried reanimation techniques and electric pacing of the posterior cricoarytenoid muscle out of 5 patients, 3 were decanulated.

Crumley RL (1990) has made a protocol for management of the injured recurrent nerve.

Some new perspectives about human laryngeal muscle: single fiber analysis and interspecies comparison were made by Wu YZ, Crumley RL, Armstrong WB and Caiozzo VJ (2000).

MedTronic has manufactured a number of prototype devices, but there are technical problems with the electrodes in the muscle. The research of electric pacing is ongoing, like in other fields the future is promising.

Genetic research of effects of denervation on cell cycle regulation of the posterior cricoartenoid muscle is also promising for new aspects of treatment as suggested by Vincent J et al. (2004).

Till prospective randomised clinical studies have been carried out we have to accept that the reinnervation studies are experimental.

The use of systemic corticosteroids and systemic antibiotics as well as antiviral treatment has to be validated.

Lx 330Deg 262.5Hz 68.3dB



SWALLOWING

The swallowing mechanism can be evaluated very specifically and gives information of further nerve involvement (Ludlow CL. 2004).

Lx 330Deg 262.5Hz 68.3dB



FOLLOW-UP

Spontaneous recovery can be expected in half of the patients. Care of complications must be made (Tucker HM).

Decannulation can be tried up to four weeks after surgery.

The follow-up in the future has to be planned ahead by statistical advice as is usual done in prospective clinical trials. Till now no suitable statistically based follow-ups have been made.



Discussion

In the future it is suggested that a prospective randomised set-up with the inclusion criteria of quality of life is made for patients with immobilisation of the vocal cords for surgery and remobilisation.

It is necessary to use evidence based clinical science to measure defined inclusion groups of patients within a protocol based on statistical calculations of difference between groups (power of study) and a local control group statistically confirmed as for amount of patients in groups, to measure difference, and a relevant follow up (Clinical Trials, Pocock SJ, 1983. John Wiley and sons).

The choice of questions to be answered are related to the inclusion criteria and eventual co-work with people that are related to funding, because the evidence based research is expensive.

Once the protocol is made, the carrying out should not wait for approval, because prospective randomised pilot studies with control groups are relevant also, and they can be the basis for clinical studies carried out later.

There does not seem to be excuses for not making evidence based documentation of surgical and non-surgical intervention.